

# Recurrent Infections May Signal Immunodeficiencies



Your immune system is constantly on the defense-fighting germs that could cause infections. Sometimes the “germ” wins but what is the difference between losing an occasional battle and having recurrent infections? How many are too many?

## Do you or your children...

- Need more than four courses of antibiotic treatment per year (in children) or more than two times per year (in adults)?
- Experience more than four new ear infections in one year after 4 years of age.
- Develop pneumonia twice over any time?
- Have more than three episodes of bacterial sinusitis in one year or the occurrence of chronic sinusitis?
- Need preventive antibiotics to decrease the number of infections?
- Develop unusually severe infections that started as common bacterial infections?

If so, these recurrent infections may be a sign of an immunodeficiency disorder.

The immune system is smart and has the ability to learn the “face” of a germ and remember it. Once your immune system has successfully battled it, most people are less susceptible to recurring infections caused by that germ.

An immunodeficiency happens when your body lacks the ability to produce antibodies which attach to germs and destroy them. Secondary Immunodeficiencies result from various conditions including cancer therapies, bone marrow transplantation and immune-modulating drugs used to treat a variety of autoimmune conditions.

Primary Immunodeficiency Diseases (PID) comprise a group of more than 100 diseases which are due to defects in the body’s immune system. These diseases affect thousands of infants, children and adults in the United States.

Most PID are inherited in our genes, so they are present at birth. Yet they may not become apparent or diagnosed until late in childhood, or even in adult life. It often takes time for a pattern of recurrent infections or other symptoms to develop before a PID is suspected.

If you think you have a PID, talk to an allergist/immunologist, a pediatrician or internist with two to three years of additional training to manage allergies, asthma and immunodeficiencies. Some PID can mimic other conditions including allergies, asthma, or eczema and vice versa, so evaluation by an allergist/immunologist is extremely helpful for reaching a diagnosis and developing an effective treatment plan.

The most common forms of immunodeficiency are often treated with infusions of antibodies called intravenous immunoglobulins (IVIG). IVIG replaces the antibodies that your body is unable to produce.

For detailed information on various forms of PID, visit [www.aaaai.org](http://www.aaaai.org).

## DID YOU KNOW?

- The first lines of defense against infection are your skin and the membranes lining the respiratory and digestive system.
- If you have year-round allergies to dust mites, pollen and mold, you may have some damage to your mucus membranes which can increase your chances of infection.

*Brought to you by your allergist/immunologist and the American Academy of Allergy, Asthma & Immunology.*  
© 2012 American Academy of Allergy, Asthma & Immunology.

[www.aaaai.org](http://www.aaaai.org)

## To the Point

People with immunodeficiency get the same kinds of infections other people get—ear infections, sinusitis and pneumonia. The difference is their infections occur more frequently, are often more severe, and have a greater risk of complications.



Henry Yeates, MD • Tammy Jacobs, MD • Evangeline Lindorf, NP  
Orem • Payson  
**801.226.3600**